# THYROID SCAR CARE GUIDE

This guide shows typical incision patterns and how silicone therapy can support healthy scar outcomes. Start silicone no sooner than 10-14 days after surgery, only when the incision is fully closed and your surgeon says it's okay.



## Standard Open "Collar" (Kocher) Incision

*CURVED, LOW IN A NECK CREASE*



## Video-Assisted/Minimally Invasive (MIVAT)

*SMALL CENTRAL INCISION*



## Extended Exposure for Cancer

*CURVED, LOW IN A NECK CREASE*




## Minimally Invasive Open (Mini-Incision) Thyroidectomy

*SMALLER CREASE OR MIDLINE CUT*


## Recommended Products

Scar is less than to two inches long:



1" x 3" Strips

Scar is more than to two inches long (trim to fit):




1" x 6" Strips

**SILAGEN + SPF30 GEL CAN ALSO BE USED IF PREFERRED**

# THYROID SCAR CARE GUIDE

This guide shows typical incision patterns and how silicone therapy can support healthy scar outcomes. Start silicone no sooner than 10-14 days after surgery, only when the incision is fully closed and your surgeon says it's okay.



**Large Goiter or Nodal Surgery**

*SUBSTERNAL/MEDIASTINAL EXTENSION*



**"Remote-Access" Robotic/ Endoscopic Approach**

*TRANSAXILLARY*



**"Remote-Access" Robotic/ Endoscopic Approach**

*BILATERAL AXILLO-BREAST (BABA)*




**"Remote-Access" Robotic/ Endoscopic Approach**

*RETROAURICULAR "FACELIFT"*


## Recommended Products

Scar is less than to two inches long:



1" x 3" Strips

Scar is more than to two inches long (trim to fit):



1" x 6" Strips

**SILAGEN + SPF30 GEL CAN ALSO BE USED IF PREFERRED**

silagen.com
@silagenscarrefinement

silagen®

MKL546V1